



GUINÉE-BISSAU (BISSAU)

Global Youth Tobacco Survey (GYTS)

Fact Sheet (Ages 13-15)



The Guinée-Bissau (Bissau) GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Guinée-Bissau (Bissau) could include in a comprehensive tobacco control program.

The Guinée-Bissau (Bissau) GYTS was a school-based survey of students in Forms 1, 2, 3, and 4 conducted in 2008.

A two-stage cluster sample design was used to produce representative data for all of Guinée-Bissau (Bissau). At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 92.0%, the class response rate was 95.6%, the student response rate was 72.5% and the overall response rate was 63.7%. A total of 1,124 13-15 year old students participated in the Guinée-Bissau (Bissau) GYTS.

Prevalence

- 5.4% of students had ever smoked cigarettes
(Male = 7.7%, Female = 3.0%)
- 10.9% currently use any tobacco product
(Male = 11.5%, Female = 10.3%)
- 5.1% currently smoke cigarettes
(Male = 7.2%, Female = 3.0%)
- 6.1% currently use other tobacco products
(Male = 4.5%, Female = 7.8%)
- 24.5% of never smokers are likely to initiate smoking next year

Cessation - Current Smokers

- 81.0% want to stop smoking
- 87.2% tried to stop smoking during the past year

Knowledge and Attitudes

- 21.6% think boys and 12.3% think girls who smoke have more friends
- 17.0% think boys and 18.4% think girls who smoke look more attractive

Access and Availability - Current Smokers

- 15.6% usually smoke at home
- 46.7% buy cigarettes in a store
- 89.0%* who bought cigarettes in a store were NOT refused purchase because of their age

* indicates cell size is less than 35

Exposure to Secondhand Smoke (SHS)

- 31.0% live in homes where others smoke in their presence
- 35.3% are around others who smoke in places outside their home
- 73.6% think smoking should be banned from public places
- 52.7% think smoke from others is harmful to them
- 18.3% have one or more parents who smoke
- 3.7% have most or all friends who smoke

Media and Advertising

- 41.2% saw anti-smoking media messages, in the past 30 days
- 32.8% saw pro-cigarette ads on billboards, in the past 30 days
- 37.2% saw pro-cigarette ads in newspapers or magazines, in the past 30 days
- 19.5% have an object with a cigarette brand logo
- 4.4% were offered free cigarettes by a tobacco company representative

School

- 33.5% had been taught in class, during the past year, about the dangers of smoking
- 29.0% had discussed in class, during the past year, reasons why people their age smoke
- 43.2% had been taught in class, during the past year, the effects of tobacco use

HIGHLIGHTS

- 10.9% of students currently use any form of tobacco; 5.1% of students currently smoke cigarettes and three in five currently use tobacco products other than cigarettes.
- SHS exposure is moderate – almost one-third of students live in homes where others smoke; more than one-third of students are exposed to smoke of others outside their home; nearly one in five students have at least one parent who smokes.
- Nearly three-quarters of students think smoking should be banned in public places.
- Two in five students saw anti-smoking media messages in the past 30 days; almost one-third of students saw pro-cigarette ads on billboards; nearly two in five students saw pro-cigarette ads in newspapers or magazines in the past 30 days.