


ADOLESCENT HEALTH IN NIGERIA




Country profile: An adolescent is a person between 10-19 years of age

0.924
185,990
199
49
41,050
22
57,226
31
2.5
39
53
18



- √ National standards for delivery of quality health services to adolescents
- ➤ No National strategies for adolescent health
- ✓ Roadmaps for accelerating HIV prevention testing, treatment and care for adolescents
- ➤ No adolescent health interventions within the HPV
- ✓ Specific target group in national policies for injury prevention among adolescents

Early marriage



Percentage of women 20-24 years married by age 18

Women 15-19 years married by age 15

M

1

Median age at 1st marriage (years)


Median age at 1st sexual intercourse (years)


Percentage of 15-19 year olds currently married or in union (%)

Pregnancy



Contraception





Demand for contraception satisfied
 Unmet need for contraception

HIV/STIs

	Comprehensive knowledge of HIV (%)			
	15-19 years		20-2	4 years
	Female	Male	Female	Male
2003	11.4		18.1	
2008	19.7	28.2	24.8	37.2
2013	22.4	29.3	26.4	38.6

	Condom use among sexually active during premarital sex (%)			
	15-19 y	ears	20-24	1 years
	Female	Male	Female	Male
2003	23.0	32.8	27.0	53.7
2008	28.2	36.2	40.9	56.3
2013	37.2	49.5	48.7	61.2

Prevalence (%) of HIV in 15-19 year olds



Prevalence (%) of STIs among 15-19 year



For further information:

Child and Adolescent health & Nutrition Programme (afrgofrhcan@who.int)
Family & Reproductive Health (FRH) cluster





ADOLESCENT HEALTH IN NIGERIA



Malnutrition


	Aged 15-19		Aged 2	0-24
	Female	Male	Female	Male
Prevalence (%) of chronically malnourished (thin, BMI<18.5)	23.1		14.4	
Prevalence (%) of overweight (BMI between 25.0-29.9)	4.9		10.9	
Prevalence (%) iron deficiency anemia (Hgb<12.0 g/dl for non- pregnant women and Hgb<11.0 g/ dl for pregnant women)				

Substance use

	Age	Female	Male
Total alcohol consumption among 15-19 year olds (litres of pure alcohol)	15-19	4.7	11.6
Prevalence cigarette or any type of	15-19	0.2	1.2
tobacco product use (%)	20-24	0.1	6.3
Prevalence chewing chat (%)	15-19		
Frevalence chewing chat (%)	20-24		



Violence







Injuries and trafic accidents

Adolescent household members injured and unable to carry out normal daily activities of >6 months (%)	
Adolescent household members injured or killed in accidental fall (%)	
Adolescent household members injured or killed in road traffic accident (%)	

27%

Sexual and gender-based violence

	by their husband or partner	aged 15-19	14.9
		aged 20-24	23.4
	Percentage of victims women of female genital mutilation (%)	aged 15-19	15.3
	` '	aged 20-24	21.7

Mental health

	Female	Male
Prevalence of childhood mental illness among 15-19 year olds (%)		
Felt seriously worthless/hopeless or wishes to be dead among15-19 year olds (%)	13	7



Immunization

Proportion of adolescent girls vaccinated for HPV to



The state of adolescent health

Birth rate per 1000 girls 15-19 year olds	122	
Girls 15-19 year olds who have given birth or are pregnant with their first child (%)	6.1	



Socioeconomic determinants of health

	Female	Male
Literacy rate among 15-19 year olds (%)	61.6	
Percent 15-19 years currently employed (%)	28.0	36.6
Information media use at least once/week among adolescents (%)	50.0	53.9
Social security health insurance among 15-19 year olds (%)	0.0	0.2

Leading causes of girls' death

Rank	Cause name	Number of deaths	Death rate/100, 000 pop
1	Lower respiratory infections	8,350	41.4
2	Diarrhoeal diseases	7,304	36.2
3	Maternal conditions	6,202	30.8
4	Meningitis	5,364	26.6
5	Sickle cell disorders and trait	5,020	24.9

Leading causes of boys' death

Rank, cause name, number of deaths and death rate, age 10-19						
Rank	Cause name	Number of deaths	Death rate/100, 000 pop			
1	Lower respiratory infections	8,800	42.1			
2	Diarrhoeal diseases	7,458	35.7			
3	Meningitis	6,311	30.2			
4	HIV/AIDS	3,291	15.8			
5	Sickle cell disorders and trait	3,232	15.5			
3	Meningitis HIV/AIDS Sickle cell disorders and	6,311				