



ADOLESCENT HEALTH IN DR CONGO



Country profile: An adolescent is a person between 10-19 years of age

2.345
78,736
34
43
17,401
22
24,966
32
3
42
60
17





✓ National standards for delivery of quality health services to adolescents


✓ National strategies for adolescent health

X No Roadmaps for accelerating HIV prevention testing, treatment and care for adolescents

X No adolescent health interventions within the HPV

X No specific target group in national policies for injury prevention among adolescents

Early marriage



M

1

Median age at 1st marriage (years)


2007

Median age at 1st sexual intercourse (years)

Percentage of 15-19 year olds currently married or in union (%) 21

Pregnancy

Percentage of 15-19 year olds who have given birth/who are pregnant with their first child



Contraception Trend in contraceptive prevalence Unmet need vs Demand satis contraception in 15-19 year olds (%) 15-19 years 50 40 satisfied Unmet need, 40 40 30 Demand 20 20 20 15 13 10 2007 2014 0 Demand for contraception satisfied

HIV/STIs

	Comp	rehensive l	knowledge o	f HIV (%)
	15-19 years 20-24 years		4 years	
	Female	Male	Female	Male
2007	13.7	18.3	16.4	23.5
2014	17.1	20.3	20.2	30.7

	Condom us	exually actives sex (%)	e during	
	15-19 ye	ears	20-24 years	
	Female	Male	Female	Male
2007	12.1	21.0	23.1	29.7
20164	22.4	26.5	30.2	29.8

Prevalence (%) of HIV in 15-19 year olds



Prevalence (%) of STIs among 15-19 year olds





For further information:

Unmet need for contraception

Child and Adolescent health & Nutrition Programme (afrgofrhcan@who.int)

2014



Adolescent Health in DR CONGO



Malnutrition

	Aged 15-19		Aged 2	20-24
	Female	Male	Female	Male
Prevalence (%) of chronically malnourished (thin, BMI<18.5)	20.8		15.4	
Prevalence (%) of overweight (BMI between 25.0-29.9)	6.2		9.5	
Prevalence (%) iron deficiency anemia (Hgb<12.0 g/dl for non- pregnant women and Hgb<11.0 g/dl for pregnant women)	40.1	37.0	39.6	18.9

Substance use

	Age	Female	Male
Total alcohol consumption among 15-19 year olds (litres of pure alcohol)	15-19	2.8	6.5
Prevalence cigarette or any type of	15-19	2.0	9.6
tobacco product use (%)	20-24	2.0	20.0
Durandaria a abandari abat (0)	15-19		
Prevalence chewing chat (%)	20-24		



Violence



Percentage of married adolescents who agree that husband is justified





75%

adolescents (%)

olds (%)

69%

•			

Mental health

Injuries and trafic accidents

Prevalence of childhood mental illness %)	
Prevalence of mental health related to alcohol, substance use, schizophrenia, seizure disorder	
and hinder disorder (%)	

Adolescent household members injured and unable

to carry out normal daily activities of >6 months (%)

Adolescent household members injured or killed in

Adolescent household members injured or killed in

accidental fall (%)

road traffic accident (%)



Sexual and gender-based violence				
% of married women who have ever experienced physical, sexual or emotional violence committed	aged 15-19	49.9		
by their husband or partner	aged 20-24	60.3		
Percentage of victims women of female genital	aged 15-19	N/A		
mutilation (%)	aged 20-24	N/A		

Immunization

Socioeconomic determinants of health

Proportion of adolescent girls vaccinated for HPV to prevent cervical cancer (%)

Literacy rate among 15-19 year olds (%)
Percent 15-19 years currently employed (%)

Information media use at least once/week among

Social security health insurance among 15-19 year



Male

90.0

37.1

48.7

0.0

Female

77.6

40.9

38.1

The state of adolescent health

Birth rate per 1000 girls 15-19 year olds	138
Girls 15-19 year olds who have given birth or are pregnant with their first child (%)	6.0

Trend in Mortality rates per 100,000 in 15-19 year olds			
2007	2014		
5.0 4.5	4.1 3.7		

Leading causes of girls' death

Rank, cause name, number of deaths and death rate, age 10-19

Rank	Cause name	Number of deaths	Death rate/100, 000 pop
1	Lower respiratory infections	2,579	29.8
2	Meningitis	1,917	22.1
3	Maternal conditions	1,770	20.4
4	Diarrhoeal diseases	1,310	15.1
5	Road injury	1,122	13.0

Leading causes of boys' death

Rank, cause name, number of deaths and death rate, age 10-19

Rank	Cause name	Number of deaths	Death rate/100, 000 pop
1	Road injury	2,757	31.5
2	Lower respiratory infections	1,888	21.6
3	Sickle cell disorders and trait	1,607	18.4
4	Drowning	1,574	18.0
5	Diarrhoeal diseases	1,526	17.5

