



# ADOLESCENT HEALTH IN MOZAMBIQUE



#### Country profile: An adolescent is a person between 10-19 years of age





 National standards for delivery of quality health services to adolescents

- National strategies for adolescent health
- × No roadmaps for accelerating HIV prevention testing, treatment and care for adolescents
- X No adolescent health interventions within the HPV vaccination
- ✓ Specific target group in national policies for injury prevention among adolescents


#### Early marriage Trend in median age of women at first marriage/sex (years) 1997 2003 2009 2011 2015 20.0 19.5 19.0 18.5 18-8 (years) 18.0 18.2 Percentage 17.5 of women 17.7 Age 20-24 years 17.5 17.0 married Ø 17.1 G by age 18 16.5 16-8 16.6 Ø G 52.9% 16.0 G 16.1 16.1 15.915.5 Women 15-19 years married Median age at 1st marriage (years) by age 15 Median age at 1st sexual intercourse (years) 129 Percentage of 15-19 year olds F Μ

Contraception

37

8

currently married or in union (%)




Demand for contraception satisfied

Unmet need for contraception

Data source: 1997-2015 DHS;UNICEF-SOWC 2017;UN DESA, Population Division. WPP 2017; The joint mission from UNICEF and WHO Africa Regional Office June 2017; http://aidsinfo.unaids.org/#; WHO-Global health observatory 2018; The World Bank 2018;WHO Global health estimates 2015, Geneva 2016.

#### Pregnancy



Women girls of 15-19 year olds who are pregnant with their first child Women girls of 15-19 years who have given birth

HIV/STIs				
Comprehensive knowledge of HIV (%)				
15-19	years	20-2	4 years	
Female	Male	Female	Male	
36.5	30.8	35.0	37.5	
27.4	48.5	33.7	56.5	1
27.7	28.0	34.1	32.9	
	15-19 Female 36.5 27.4	IS-19 years   Female Male   36.5 30.8   27.4 48.5	Comprehensive knowledge of   15-19 years 20-2   Female Male Female   36.5 30.8 35.0   27.4 48.5 33.7	Comprehensive knowledge of HIV (%)   15-19 years 20-24 years   Female Male Female Male   36.5 30.8 35.0 37.5   27.4 48.5 33.7 56.5

	Condom use among sexually active during premarital sex (%)				
	15-19 y	ears	20-24	1 years	
	Female	Male	Female	Male	
2009	44.5	37.1	46.7	56.1	
2011	43.7	41.1	49.6	54.1	
2015	52.3	40.4	61.6	58.0	


Prevalence (%) of HIV in 15-19 vear olds



Prevalence (%) of STIs among 15-19 year olds



Africa



0

Child and Adolescent health & Nutrition Programme (afrgofrhcan@who.int) Family & Reproductive Health (FRH) cluster



# **ADOLESCENT HEALTH IN MOZAMBIQUE**



### **Malnutrition**

	Aged 15-19		Aged 20-24	
	Female	Male	Female	Male
Prevalence (%) of chronically malnourished (thin, BMI<18.5)	14.5		6.4	
Prevalence (%) of overweight (BMI between 25.0-29.9)	6.2		9.2	
Prevalence (%) iron deficiency anemia (Hgb<12.0 g/dl for non- pregnant women and Hgb<11.0 g/dl for pregnant women)	54.9		53.3	

#### Violence



Percentage of married adolescents who agree that husband is justified

20%



· · · · ·

### Sexual and gender-based violence

private sexual of emotional violence commuted	aged 15-19	14.7
by their husband or partner	aged 20-24	21.4
Percentage of victims women of female genital mutilation (%)	aged 15-19	N/A
	aged 20-24	N/A

#### Immunization

Proportion of adolescent girls vaccinated for HPV to prevent cervical cancer (%)

## Socioeconomic determinants of health

	Female	Male
Literacy rate among 15-19 year olds (%)	58.3	78.3
Percent 15-19 years currently employed (%)	21.1	39.3
Information media use at least once/week among adolescents (%)	56.6	73.1
Social security health insurance among 15-19 year olds (%)		

## Leading causes of girls' death

Rank, cause name, number of deaths and death rate, age 10-19					
Rank	Cause name	Number of deaths	Death rate/100, 000 pop		
1	HIV/AIDS	1,096	32.9		
2	Lower respiratory infections	811	24.4		
3	Diarrhoeal diseases	741	22.3		
4	Meningitis	531	16.0		
5	Maternal conditions	446	13.4		

Data source: 1997-2015 DHS;UNICEF-SOWC 2017;UN DESA, Population Division. WPP 2017; The joint mission from UNICEF and WHO Africa Regional Office June 2017; http://aidsinfo.unaids.org/#; WHO-Global health observatory 2018; The World Bank 2018;WHO Global health estimates 2015, Geneva 2016.

# Substance use

	Age	Female	Male	
Total alcohol consumption among 15-19 year olds (litres of pure alcohol)	15-19	3.2	5.4	
Prevalence cigarette or any type of	15-19	0.3	4.3	
tobacco product use (%)	20-24	0.9	17.0	
Dravelance showing shot (%)	15-19			
Prevalence chewing chat (%)	20-24			

# Injuries and trafic accidents

Adolescent household members injured and unable to carry out normal daily activities of >6 months (%) Adolescent household members injured or killed in accidental fall (%) Adolescent household members injured or killed in

road traffic accident (%)

## Mental health

Prevalence of childhood mental illness %) Prevalence of mental health related to alcohol, substance use, schizophrenia, seizure disorder

and bipolar disorder (%)




. . .

. . .

. . .

### The state of adolescent health



### Leading causes of boys' death

Rank, cause name, number of deaths and death rate, age 10-19					
Rank	Cause name	Number of deaths	Death rate/100, 000 pop		
1	HIV/AIDS	999	30.1		
2	Diarrhoeal diseases	894	26.9		
3	Road injury	648	19.5		
4	Lower respiratory infections	585	17.6		
5	Meningitis	474	14.3		

Child and Adolescent health & Nutrition Programme (afrgofrhcan@who.int) Family & Reproductive Health (FRH) cluster



For further information: