


ADOLESCENT HEALTH IN KENYA 2018




Country profile: An adolescent is a person between 10-19 years of age

T ()	
Total surface area (million square km)	0.5804
Total population (000's)	48,461
Population density (persons/ square km)	83
Urbanized population (%)	26
Adolescent population (000's)	11.1
Adolescent as % of total population	22.8
Young people (10-24 years) (000's)	14,606
Young people as % of total population	31.7
Annual average growth rate of population (%)	2.3
Crude birth rate (births/1,000 population)	31.3
Life expectancy at birth (years)	67.0
Median age in total population (years)	18.9
Birth rate per 1000 girls 15-19 years olds	96








adolescent health


 Roadmaps for accelerating HIV prevention testing, treatment and care for adolescents

✓ Adolescent health interventions within the HPV vaccination


X No Specific target group in national policies for injury prevention among adolescents



Contraception



Pregnancy



Girls of 15-19 year olds who are pregnant with their first child Women girls of 15-19 years who have given birth

				3
	nowledge o	of HIV		
	15-19	15-19 years		years
	Female	Male	Female	Male
2003	27.3	35.9	40.9	61.4
2008	42.0	51.2	51.4	60.5
2014	49.0	57.7	59.6	70.9

	Condom	ve during			
	15-19 years		20-24	Years	
	Male	Female	Male	Female	
1998	37.8	16.1	49.1	18.3	
2003	41.1	24.7	52.3	30.5	
2008	54.9	41.8	69.4	38.9	
2014	66.3	55.6	79.0	64.8	

HIV/STIs

Prevalence (%) of HIV in 15-19 year olds

†

Prevalence (%) of STIs among 15-19 year



World Health Organization

Data source: 1989-2013 DHS; UNICEF-SOWC 2017; UN DESA, Population Division. WPP 2017; The joint mission from UNICEF and WHO Africa Regional Office June 2017; http://aidsinfo.unaids.org/#; WHO-Global health observatory 2018; The World Bank 2018; WHO Global health estimates 2015, Geneva 2016.

For further information:

Child and Adolescent health & Nutrition Programme (afrgofrhcan@who.int) Family & Reproductive Health (FRH) cluster



ADOLESCENT HEALTH IN KENYA 2018



. . .

. . .

Malnutrition

		15-19 years		20-24 years	
		Female	Male	Female	Male
Prevalence (%) of chronically malnourished (thin, BMI<18.5)	1998	17.4		10.5	
	2003	20.4		9.3	
	2008	19.0		11.3	
	2014	16.6		9.6	
Prevalence (%) of overweight (BMI between 25.0-29.9)	1998	8.9		9.5	
	2003	7.5		14.6	
	2008	7.2		13.2	
	2014	10.6		18.0	

Violence



Percentage of married adolescents who agree that husband is justified in beating his wife

45%

rbeating	nis wie			
	37%			
				Female
on who l		15-19 years	2008	33.6

	15-19 years	2008	33.6
% of married women who have ever experienced physical or sexual or emotional violence		2014	30.7
committed by their husband or partner	20-24 years	2008	40.7
		2014	41.5

Immunization

Proportion of adolescent girls vaccinated for HPV to prevent cervical cancer (%)

Socioeconomic determinants of health

		Female	Male
Literacy rate among 15-19 year olds (%)			
		22.3	16.8
Percent 15-19 year currently employed (%)	2003	28.6	25.5
	2008	19.3	59.9
		18.8	34.7
Information media use at least once/week among adolescents (%)	2016	74.8	83.8
Social security health insurance among 15-19 year olds (%)		0.4	0.5
		5.6	7.2

Leading causes of girls' death

Rank, cause name, number of deaths and death rate, age 10-19				
Rank	Cause name	Number of deaths	Death rate/100, 000 pop	
1	HIV/AIDS	1,326.07	24.1	
2	Diarrhoeal diseases	1,118.84	20.3	
3	Malaria	896.27	16.3	
4	Lower respiratory infections	685.96	12.5	
5	Maternal conditions	603.09	11.0	

Data source: 1989-2013 DHS;UNICEF-SOWC 2017;UN DESA, Population Division. WPP 2017; The joint mission from UNICEF and WHO Africa Regional Office June 2017 ; http://aidsinfo.unaids.org/#; WHO-Global health observatory 2018; The World Bank 2018;WHO Global health estimates 2015, Geneva 2016. ...: no data; n/a: not applicable

Substance use

			Female	Male	1005
Total alcohol consumption among 15-19 year olds (litres of pure alcohol)	15-19 years	2010	2.8	9.7	
	15-19 years	2003	1.0	5.6	
	:	2008	0.4	3.0	
Prevalence cigarette or any other		2014	0.2	2.2	
tobacco product use (%)		2003	1.2	21.9	
		2008	1.0	16.6	
		2014	1.0	12.1	

Injuries and trafic accidents

Adolescent household members injured and unable to carry out normal daily activities of >6 months (%) Adolescent household members injured or killed in accidental fall (%) Adolescent household members injured or killed in

road traffic accident (%)

Mental health

	Female	Male	
Prevalence of childhood mental illness among 15-19 year olds (%)			N
Felt seriously worthless/hopeless or wishes to be dead among15-19 year olds (%)			

Adolescent Mortality



Leading causes of boys' death

Rank, cause name, number of deaths and death rate, age 10-19

	, , ,		, 0
Rank	Cause name	Number of deaths	Death rate/100, 000 pop
1	HIV/AIDS	1,642.00	29.5
2	Diarrhoeal diseases	1,541.80	27.7
3	Road injury	772.56	13.9
4	Malaria	686.46	12.3
5	Meningitis	676.77	12.2

For further information:

F

Child and Adolescent health & Nutrition Programme (afrgofrhcan@who.int) Family & Reproductive Health (FRH) cluster

