



ADOLESCENT HEALTH IN CAR 2018



Country profile: An adolescent is a person between 10-19 years of age



Early marriage






currently married or in union (%)



Pregnancy



Girls of 15-19 year olds who are pregnant with their first child Women girls of 15-19 years who have given birth

HIV/STIs							
	Comp	Prevalence (%) of HIV in 15-19 year olds					
Fe		emale	М	ale			
		N	o Data		\mathbf{P}		
Condom use among sexually active during premarital sex (%)					Prevalence (%) of STIs among 15-19 year		
		years		Years	ÅÅ		
	Female	Male	Female	Male	22		

World Health

Organization

FICE FOR Africa

17.0

Data source: 1994 DHS;UNICEF-SOWC 2017;UN DESA, Population Division. WPP 2017; The joint mission from UNICEF and WHO Africa Regional Office June 2017; http://aidsinfo.unaids.org/#; WHO-Global health observatory 2018; The World Bank 2018;WHO Global health estimates 2015, Geneva 2016.

For further information:

1994

Child and Adolescent health & Nutrition Programme (afrgofrhcan@who.int) Family & Reproductive Health (FRH) cluster

15.1



ADOLESCENT HEALTH IN CAR 2018



Malnutrition					
15-19 years 20-24 years					years
		Female	Male	Female	Male
Prevalence (%) of chronically malnourished (thin, BMI<18.5)	1994	15.2		14.8	
Prevalence (%) of overweight (BMI between 25.0-29.9)	1994	3.2		3.9	

Violence



79%

Percentage of married adolescents who agree that husband is justified in beating his wife

Ś.

-

Substance use

			Female	Male	19 SA
Total alcohol consumption among 15-19 year olds (litres of pure alcohol)	15-19 years	2010	4.5	8.7	
Prevalence cigarette or any other tobacco product use (%)	15-19 years				

Injuries and trafic accidents

Adolescent household members injured and unable
to carry out normal daily activities of >6 months (%)...Adolescent household members injured or killed in
accidental fall (%)...Adolescent household members injured or killed in
road traffic accident (%)...

Mental health

	Female	Male	
Prevalence of childhood mental illness among 15-19 year olds (%)			N
Felt seriously worthless/hopeless or wishes to be dead among15-19 year olds (%)			

Immunization

Proportion of adolescent girls vaccinated for HPV to prevent cervical cancer (%)


Socioeconomic determinants of health

			Male
Literacy rate among 15-19 year olds (%)			
Percent 15-19 year currently employed (%)	1994	54.4	48.7
Information media use at least once/week among adolescents (%)	2016		
Social security health insurance among 15-19 year olds (%)			

Leading causes of girls' death

Rank, cause name, number of deaths and death rate, age 10-19						
Rank	Cause name	Number of deaths	Death rate/100, 000 pop			
1	HIV/AIDS	192	34.1			
2	Lower respiratory infections	184	32.8			
3	Maternal conditions	158	28.2			
4	Diarrhoeal diseases	155	27.5			
5	Meningitis	145	25.7			

Data source: 1994 DHS;UNICEF-SOWC 2017;UN DESA, Population Division. WPP 2017; The joint mission from UNICEF and WHO Africa Regional Office June 2017; http://aidsinfo.unaids.org/#; WHO-Global health observatory 2018; The World Bank 2018;WHO Global health estimates 2015, Geneva 2016. ...: no data; n/a: not applicable Adolescent Mortality



Leading causes of boys' death

Rank, cause name, number of deaths and death rate, age 10-19

Name, cause name, number of deaths and death rate, age 10-13						
Rank	Cause name	Number of deaths	Death rate/100, 000 pop			
1	Diarrhoeal diseases	193	34.7			
2	HIV/AIDS	185	33.3			
3	Lower respiratory infections	164	29.5			
4	Drowning	136	24.5			
5	Meningitis	116	20.9			

For further information:

Child and Adolescent health & Nutrition Programme (afrgofrhcan@who.int) Family & Reproductive Health (FRH) cluster

