


ADOLESCENT HEALTH IN ZAMBIA 2018

Country profile: An adolescent is a person between 10-19 years of age

Total surface area (million sq. km)	0.7526	
Total population (000's)	16,591.3900	0
Population density (persons/sq km)	21.6583	7
Urbanized population (%)	41.3790	
Adolescent population (000's)	3,955.8010	
Adolescent as % of total population	23.8425	
Young people (10-24 years) (000's)	5,355.3410	1
Young people as % of total population	33.0316	
Annual average growth rate of population (%)	2.8882	
Crude birth rate (births per 1,000 population)	38.1310	
Life expectancy at birth (years)	61.8740	1
Median age of the total population (years)	17.0510	
Birth rate per 1000 girls 15-19 years olds	145.0000	

Early marriage




Percentage of 15-19 year olds currently married or in union (%)




F


17

31%





Pregnancy



Girls of 15-19 year olds who are pregnant with their first child Women girls of 15-19 years who have given birth

	HIV/STIs				
	Comprehensive knowledge of HIV				
	15-19	years	s 20-24 years		
	Female	Male	Female	Male	
2002	32.2		37.2		
2007	32.3	35.0	36.0	39.5	
2014	38.9	42.3	44.7	53.0	
	Condom use among sexually active during premarital sex (%)				
	15 19 years 2			voare	

	15-19 years		20-24	years
	Male	Female	Male	Female
2002	32.9	29.6	51.2	38.4
2007	41.7	36.0	52.9	44.4
2014	42.5	36.5	55.5	44.7

Prevalence (%)

of HIV in 15-19 vear olds



Prevalence (%) of STIs among 15-19 year



ICE FOR Africa

World Health

Organization

For further information: Child and Adolescent health & Nutrition Programme (afrgofrhcan@who.int)

Family & Reproductive Health (FRH) cluster

Data source: 1992-2016 DHS:UNICEF-SOWC 2017;UN DESA, Population Division. WPP 2017; The joint mission from UNICEF and WHO Africa Regional Office June 2017; http://aidsinfo.unaids.org/#; WHO-Global health observatory 2018; The World Bank 2018;WHO Global health estimates 2015, Geneva 2016.



ADOLESCENT HEALTH IN ZAMBIA 2018



Malnutrition

		15-19 years		20-24 years	
		Female	Male	Female	Male
	1996	16.8		14.6	
Prevalence (%) of chronically malnourished	2001	15.8		11.0	
(thin, BMI<18.5)	2006	13.9		9.4	
	2012	13.2		7.4	
	1996	2.0		4.2	
Prevalence (%) of overweight (BMI between	2001	5.9		11.2	
25.0-29.9)	2006	6.0		9.3	
	2012	9.2		13.5	
Prevalence (%) of iron deficiency anemia (Hgb<12.0 g/dl for non-pregnant women and Hgb<11.0 g/dl for pregnant women)	2001	65.2		65.4	
	2006	59.2		63.1	
	2012	41.6		41.0	

Violence



Percentage of married adolescents who agree that husband is justified in beating his wife



· · · · ·

			Female
% of married women who have ever experienced physical or sexual or emotional violence committed by their husband or partner	15-19 years	2013	40.1
	20-24 years	2013	47.9
Prevalence of victims women of femal genital mutilation (%)	15-19 years	2014	
	20-24 years	2014	

Immunization

Proportion of adolescent girls vaccinated for HPV to prevent cervical cancer (%)

Socioeconomic determinants of health

		Female	Male
Literacy rate among 15-19 year olds (%)	2014	78.8	82.3
Percent 15-19 year currently employed (%)	2014	19.4	36.5
Information media use at least once/week among adolescents (%)	2016	69.1	75.3
Social security health insurance among 15-19 year olds (%)	2013	0.0	0.0

Leading causes of girls' death

Rank, cause name, number of deaths and death rate, age 10-19							
Rank	Cause name	Number of deaths	Death rate/100, 000 pop				
1	HIV/AIDS	639	32.4				
2	Malaria	469	23.8				
3	Meningitis	273	13.9				
4	Diarrhoeal diseases	269	13.7				
5	Lower respiratory infections	231	11.7				

Data source: 1992-2016 DHS;UNICEF-SOWC 2017;UN DESA, Population Division. WPP 2017; The joint mission from UNICEF and WHO Africa Regional Office June 2017 ; http://aidsinfo.unaids.org/#; WHO-Global health observatory 2018; The World Bank 2018;WHO Global health estimates 2015, Geneva 2016. ...: no data; n/a: not applicable

Substance use

			Female	Male	1 1 S
Total alcohol consumption among 15-19 year olds (litres of pure alcohol)	15-19 years	2010	2.8	6.4	
Prevalence cigarette or any other tobacco product use (%)	15-19 years	2013	0.1	3.3	
	20-24 years	2013	0.5	13.8	

Injuries and trafic accidents

Adolescent household members injured and unable
to carry out normal daily activities of >6 months (%)...Adolescent household members injured or killed in
accidental fall (%)...Adolescent household members injured or killed in
......

road traffic accident (%)

Mental health

	Female	Male	
Prevalence of childhood mental illness among 15-19 year olds (%)			N
Felt seriously worthless/hopeless or wishes to be dead among15-19 year olds (%)			

Adolescent Mortality

Trend in Mortality rates per 100,000 in 15-19 year olds 1996 2002 2007 2014 4.3 1.4 4.4 4.1 1.9 1.9 1.9 3.4 4.4 4.1 1.9 1.9 1.9 1.9 2.4 2.1 1.9 1.9 1.9 1.9 1.9 1.9 4.3 1.4 4.1 1.1 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9 1.9

Leading causes of boys' death

Rank, cause name, number of deaths and death rate, age 10-19						
Rank	Cause name	Number of deaths	Death rate/100, 000 pop			
1	HIV/AIDS	717	36.1			
2	Malaria	367	18.5			
3	Diarrhoeal diseases	338	17.0			
4	Meningitis	309	15.6			
5	Lower respiratory infections	270	13.6			

For further information: Child and Adolescent health & Nutrition Programme (afrgofrhcan@who.int)

Child and Adolescent health & Nutrition Programme (afrgofrhcan@who.int) Family & Reproductive Health (FRH) cluster

