


# FAMILY PLANNING HEALTH PROFILE **WHO AFRICAN REGION 2019**

General statistics			Summary FP statistics		
Total population (000's)	2019	1,091,759	Demand satisfied (%)	2019	56
Total fertility rate (births per women)	1988-2018	5	mCPR (%)	2019	24
Adolescent Birth Rate (births per 1000 adolescent females)	2015-2020	102	Unintended pregnancies (000's)	2019	22,689
IMR (deaths per 1000 live births)	2018	42	Unmet need (%)	2019	20
MMR (deaths per 1000 live births)	2017	525	Unsafe abortions averted (000's)	2019	5,377



### mCPR by place of residence (2010-2018)Urban Rural 29% 16% Availability of service delivery points (SDP) (2018) Primary SDPs with at least 3 89% modern methods

Secondary/tertiary SDPs with at

least 5 modern methods

#### (2018)12% Long-acting Implant and 21% permanent.. IUD 399 Short-term Emergency .. methods 11% Injectable Pill 13%

	(2018)				
	Long-acting and permanent	Sterilization	<b>†</b> 14%		
me %	methods	Sterilization.	<b>†</b> 15%		
	Short-term methods	Condoms	<b>i</b> 34%		
		Condoms.	<b>†</b> 8%		

## **Definition of Indicators**

- 1. mCPR: % of women of reproductive age who are using (or whose partner is using) a modern contraceptive method at a particular point in
- 2. unmet need: % of fecund women of reproductive age who want no more children or to postpone having the next child, but are not using a contraceptive method, plus women who are currently using a traditional method of family planning.

  3. demand satisfied: percentage of women (or their partners) who desire either to have no additional children or to postpone the next child and
- who are currently using a modern contraceptive method.

  4. Unintended pregnancies: Nb of pregnancies that occurred at a time when women (and their partners) either did not want additional children or wanted to delay the next birth.
- 5. unsafe abortion: Nb of unsafe abortions that did not occur during a specified reference period as a result of the protection provided by modern contraceptive use during the reference period.
- 6.Facilities stocked out: % of facilities stocked out of each type of contraceptive offered, on the day of assessment.

79%

- 7. Method availability: % of primary (resp. seconday/tertiary) service delivery points with at least 3 (resp. 5) modern methods of contraception
- 8. Total fertility rate: Average nb of children a hypothetical cohort of women would have at the end of their reproductive period if they were subject during their whole lives to the fertility rates of a given period and if they were not subject to mortality. 9. Adolescent birth rate (ABR): Nb of births to adolescent females, aged 15-19 occurring during a given reference period per 1,000 adolescent
- females. Maternal mortality ratio (MMR): Annual nb of female deaths from any cause related to or aggravated by pregnancy or its management per
- 100 000 live births.
- 11. Infant mortality rate (IMR): probability of dying between birth and age 1 per 1000 live births.

## References

From UN DESA population division 2020 Estimates and projections of family planning indicators

(estimates for Sub-Saharan Africa) Indicators

From WHO Global Observatory

\*Disaggregated indicators - median of country values

Indicators 4 to 8

Median of country values

Indicator 9 WHO Global Observatory

Indicator 10

From UN Inter-agency Group for Mortality

Estimation (MMEIG), 2016

Indicator 1

Median of country values from UN Inter-agency Group for Child Mortality (IGME), 2016

