

FAMILY PLANNING HEALTH PROFILE KENYA 2019



		Conorol	atatiatiaa						Sumr	nonu		otatia	tion				
			statistics						Sum	i ai y	FF 8	statis	ucs				
Total population (000's) 2019			2019		Demand satisfied (%)					2	2019				78		
women)			2014		mCPR (%)					2	2019			45			
Adolescent Birth Rate (births per 1000 adolescent females) 207			2014		96	Unintended pregnancies				(000	's) 2	2019			2,356		
IMR (deaths per 1000 live births) 2018			2018		Unmet need (%)					2	2019				1		
MMR (deaths per 1000 live births) 2017			s) 2017		Unsafe abortions ave				d (000)'s) 2	2019				519		
Key Family Planning Indicators by 5-year Age group (2018)				mCPR by economic status			tus (2	ıs (2018) Treno				d of key FP indicators (2012-2019)					
15-19	57	30	23	Lowest			6) 48		80	75	76	77	78 7	87	8 78	7	
20-24	81	60	14						70 60								
25-29	82	63	14	Second				67 🌖	05 pt		40	42 4	44 4	44		-4	
30-34	81	65	15	Middle				6 3	50 Bercentage 30 30	00-					45		
35-39	81 79	67	16						20	23	22	20	18 1			1	
40-44	79	59	16	Fourth				9 63	10 0				18 1	81	8 17		
45-49	2018	43 2018	12 2018	Highest				6 2		2012	2013	2014	2015		2018	2019	
Demand s	atisfied (%) mCPR (et need (%)						Unmet ne	ed (%)	m	CPR (%)		Der	nand satisf	ied (%)	
	R by place o			Facilities	s stocked (20		by me	ethod		acili	ities s		ed oı 018)	ıt by	metl	nod	
Urban Rural 61% 61%			Long-acting	<u>`</u>						-acting		Sterilization		†			
Availability of service delivery (SDP) (2018)		ry points	and permanent				and perman methods				rilizati	on.	İ				
Primary SDPs with at least 3 97%			07%	Short-term methods	e			Sh	Short-term		Condoms			No Data			
modern methods Secondary/tertiary SDPs with at				mounouo					methods			Condoms.					
	modern meth		58%		Pill										ï	1	
				n of Indicate								R	efere	ence	s		
me. unmet ne contraceptiv demand s demand s Unintend or wanted to unsafe al nodern con Facilities s Method a Method a Total ferti- ubject duri	% of women of repro- sed: % of fecund wo ve method, plus wo satisfied: percentag rently using a mode ded pregnancies: Nb o delay the next birt bortion: Nb of unsaf traceptive use duri stocked out: % of fa availability: % of piri availability: % of piri iday of assessmen ility rate:Average ht ing their whole lives ent birth rate (ABR):	omen of reproductive men who are curre e of women (or the ern contraceptive n of pregnancies th h. e abortions that did g the reference pe cilities stocked out mary (resp. second t. o of children a hypo to the fertility rates	ve age who want r ntly using a traditi ir partners) who d nethod. at occurred at a tin d not occur during eriod. to f each type of cc ay/tertiary) service othetical cohort of s of a given period	no more children or onal method of fam lesire either to have me when women (a a specified referen ontraceptive offerec e delivery points wil women would have and if they were no	to postpone h ily planning. no additional and their partne ice period as a d, on the day o th at least 3 (re e at the end of ot subject to m	aving the r children o ers) either result of t f assessm esp. 5) mo their repro ortality.	next chil r to post did not v he prote ent. dern me ductive	d, but are pone the vant addi ction prov thods of o period if t	not using next child tional child vided by contracepti hey were	a F n and o A ren <u>S</u> Ir M e Ir on V F eent E	ndicators rom FP2/ ational su ther natic ther natic ther natic ther natic ther natic ther natic ther nation naticator 8 NCAH D Stimate) ndicator 9 VHO Glob NICS,RHS ndicator 1 rom UN 1 stimation ndicator 1	020 2019 urveys (E onal surv /progres Data Porta Data Porta Data Obse S and oth 0 Inter-age o (MMEIC	DHS,PM, eys). ss.family al (UN p ervatory her natio	A2020, plannir opulatio (DHS,F nal sum	MICS,RH 192020.c on divisio 19MA2020 reys).	HS and org/re- n: 2019	

100 000 live births. 11. Infant mortality rate (IMR): probability of dying between birth and age 1 per 1000 live births.

For further information: Reproductive Maternal Health & Ageing(RMH) Programme (afrgofrhrwh@who.int), Universal Health Coverage/Life course Cluster

(IGME), 2016

