



ABORTION HEALTH PROFILE

CONGO 2019

General Statistics		
Total population (000's), 2019	5,381	
Total female population (15 - 49 years) (000's), 2019	1,295	
Unintended pregnancies (000's), 2019	115	
Adolescent birth rate (per 1000 women aged 15-19 years), 2013	111	
Proportion of women who make their own informed decisions regarding sexual relations (% of women aged 15-49 years), 2012	71	
Total fertility rate 15-49 (births per woman), 2011	5	

Legal status of abortion		
	Not specified	No
On request		×
For economic or social reasons		×
In cases of foetal impairment		×
In cases of incest		×
In cases of intellectual/cognitive disability of woman		×
In cases of rape		×
To preserve a woman's health		×
To preserve a woman's mental health		×
To preserve a woman's physical health		×
To save a woman's life		×
Other	0	





5. Annual average rate of reduction: 2010-2017 vs 2010-2030

Current AARR in MMR 4.2% Expected AARR to reach MMR target 2030

averted due to modern contraceptive use 24 2124 24 24 19 20 18 18 19 17 19 15 16 10 5 6 5 0 2012 2013 2014 2015 2016 2017 2018

6. FP indicators and unsafe abortions

7. Available in National Essential Medicines List - (2013)

Misoprostol X Mifepristone

8. Post-abortion contraceptive acceptance

6.4%

No data

References

- 1: United Nations, Department of Economic and Social Affairs, Population Division (2019). World Population Prospects 2019, custom data acquired via website in July 2020; DHS and national surveys compiled in WHO Global Observatory and UNFPA global databases accessed in July 2020; ICF, 2015. The DHS Program STATcompiler. Funded by USAID. Available from: http://www.statcompiler.com. Accessed July 2020
- 2: Global Abortion Policy Database (https://abortion-policies.srhr.org/) accessed in July 2020 3: Global Burden of Disease Study 2017 (GBD 2017) Results. Seattle, United States: Institute for Health Metrics and Evaluation (IHMF) 2018

Available from http://ghdx.healthdata.org/gbd-results-tool

Misoprostol + Mifepristone combination

- 4: Trends in maternal mortality 2000 to 2017: estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations Population Division
- 6: From FP2020 2019 Progress Report, most recent national surveys (DHS,PMA2020, MICS,RHS and other national surveys).
- lanning2020.org/resources
- 7: Global Abortion Policy Database (https://abortion-policies.srhr.org/) accessed in July 2020.

For further information:

Contraceptive prevalence - modern methods (%)

■ Demand satisfied for modern contraception (%)

Unmet need for family planning (%)

■ Unsafe abortions averted (000's)

Reproductive Maternal Health & Ageing(RMH) Programme (afrgofrhrwh@who.int), Universal Health Coverage/Life course Cluster



Definitions and abbreviations

Unsafe abortions averted due to modern contraceptive use: Number of unsafe abortions that did not occur during a specified reference period as a result of the protection provided by modern contraceptive use during the reference period. Contraceptive prevalence - Modern Methods: Percentage of women of reproductive age who are using (or whose partner is

using) a modern contraceptive method at a particular point in time

Unmet need: percentage of fecund women of reproductive age who want no more children or to postpone having the next child, but are not using a contraceptive method, plus women who are currently using a traditional method of family planning FP - Family planning, MMR - Maternal Mortality Ratio