

Mpox Incident Management Team Ministry of Health, Kampala, Uganda

Email: info@health.go.ug Hotline: 0800-100-066 Website: www.health.go.ug Date: September 2, 2025

National Mpox Situation Report

1 Summary

Total Cases	New Last 24h	Number of Deaths	Case Fatality Rate (%)
8,001	15	50	0.62

Key Points:

- 1. Epidemic remains active. Todate 82% (120/146) of districts resported at least one case since the beginning of the outbreak and 27%(40/146) reporting new cases in the past 21 days.
- 2. Overall decline. There has been an overall decline in the weekly incident cases reported. For the first time since October 2024, fewer than 100 incident cases have been reported each week for six consecutive epidemiological weeks
- 3. **Flattening curve.** There is an apparent flattenting of the curve as indicated in the cumulative curve in **figure 2.7** over the last 30 days.
- 4. High-risk mortality observed. Cumaulatively 50 deaths have occurred among confirmed cases, nearly half (46%) of whom were co-infected with HIV.
- 5. Disproportionate deaths in males and females An equal number of deaths occurred among males and females (25 each). Among the male deaths, 72% (18/25) were below 30 years of age, whereas among the female deaths, 68% (17/25) were aged above 30 years.

Weekly and Cumulative Confirmed Mpox Cases ¹					
Year	Epi Week	Weekly Confirmed Cases	Cumulative Confirmed Cases	Percent Change (%)	
2025	27	173	7404	2.39	
2025	28	108	7512	1.46	
2025	29	74	7586	0.99	
2025	30	124	7710	1.63	
2025	31	83	7793	1.08	
2025	32	71	7864	0.91	
2025	33	10	7874	0.13	
2025	34	31	7905	0.39	
2025	35	81	7986	1.02	
2025	36	15	8001	0.19	

¹ This is data for the last 10 Epiweeks. % Week-on-Week Change

2 Surveillance and Epidemiology

2.1 Summary Table of Incident Cases by Age and Sex

Cases by Age Group and Gender

Distribution across demographics

Age Group	Total Cases	% of Cases ¹	Males	% of All Males ²	Females	% of All Females ²
0-4	351	4.6%	202	4.6%	149	4.6%
5-9	327	4.3%	184	4.2%	143	4.4%
10-14	230	3.0%	125	2.9%	105	3.3%
15-19	704	9.3%	334	7.7%	370	11.5%
20-24	1,263	16.7%	561	12.9%	702	21.8%
25-29	1,582	20.9%	867	19.9%	715	22.2%
30-34	1,181	15.6%	721	16.5%	460	14.3%
35-39	905	11.9%	611	14.0%	294	9.1%
40-44	518	6.8%	363	8.3%	155	4.8%
45-49	310	4.1%	230	5.3%	80	2.5%
50+	206	2.7%	164	3.8%	42	1.3%
_	7,577 ³	100.0%	4,362	100.0%	3,215	100.0%


¹ Percentage of all cases within each age group


2.2 Summary Table of Incident Cases by the Top 10 Districts


Top 10 Districts Summary					
District	Total Cases	Total Deaths	Cases Last Epiweek	Deaths Last Epiweek	% Change in Cases
Kampala District	2,718	16	2	0	100.0%
Wakiso District	956	8	0	0	-100.0%
Mbarara City	698	5	0	0	-100.0%
⁄lasaka City	353	3	0	0	-100.0%
Mukono District	289	2	0	0	-100.0%
loima City	212	2	2	0	-33.3%
yantonde District	143	0	0	0	NA
Nakasongola District	128	1	0	0	NA
uwero District	112	0	0	0	NA
Mubende District	109	0	0	0	NA

¹ The percentage change in cases is calculated by comparing the number of cases in the most recent epidemiological week to the previous epidemiological week.

2.3 Proprotion of Affected Districts









² Percentage of all cases within each gender category

 $^{^{^{3}}\,\}mathrm{Total}$ cases less due to some missing age information

Overall Distribution of Incident Cases by Time 2.4




Epicurves of the Top 10 Districts 2.5




Overall Distribution of Incident Cases by Time 2.6


2.7 Weekly Cumulative Cases




n = 7966 from National level; Case reported range from Tue 09 Jul 2024 to Mon 01 Sep 2025 0 cases missing date of onset and not shown


2.8 Overall Distribution of Incident Cases by Age and Sex







2.9 Overall Distribution of Deaths by Place and Sex - Age







2.10 Distribution of Deaths by HIV Status







2.11 Overall Attack Rates by Age







2.12 Distribution of Incident Cases by Place(Absolute Numbers)







2.13 Attack Rates per 100K Population by Place





2.14 Attack Rates per 100K Population by Place and Top 10 Districts







CASE DEFINITION

COMMUNITY CASE DEFINITION

 An Illness with signs of skin rash, fever, headache, muscle aches, back pain, and body weakness.

SUSPECTED CASE

 Any person with skin rash on any part of tthe body not expained by any locally relevant common cause of papular or vesicular rash, mucosal lesions or swollen lymph nodes with acute onset of fever (>38.5 °C), hadache, myalgia (muscle pain/body ache), back pain, profound weakness, or fatigue.

PROBABLE CASE

 A suspect case with history of contact to probable or confirmed Mpox case 21 days before the onset of signs pr symptoms

OR

 Has an epidemiological link to a probable or confirmed case of Mpox in the 21 days before symptom onset including multiple and/or casual sexual partners in the 21 days before symptom onset: has a positive test result for orthopoxviral infection (e.g., OPXV-specific PCR without MPXV-specific PCR or sequencing)



CONFIRMED CASE

 A person with laboratory confirmed MPXV infection by detection of Unique sequences of Viral DNA by real-time polymerase chain reaction (RT-PCR) and/or sequence

Contact Us

Twitter: @MinofHealthUG Website: https://health.go.ug/ Talk to Us: 0800-100-066

Write to Us: helpdesk@health.go.ug

