



# National Mpox Situation Report

## 1 Summary

Total Cases	New Last 24h	Number of Deaths	Case Fatality Rate (%)
8,041	3	50	0.62

### Key Points:

- Epidemic remains active.** To date **82% (120/146)** of districts reported at least one case since the beginning of the outbreak and **27% (40/146)** reporting new cases in the past **21 days**.
- Overall decline.** There has been an overall decline in the weekly incident cases reported. For the first time since October 2024, fewer than 100 incident cases have been reported each week for **seven consecutive epidemiological weeks**.
- Flattening curve.** There is an apparent flattening of the curve as indicated in the cumulative curve in **figure 2.7** over the last **40 days**.
- High-risk mortality observed.** Cumulatively **50 deaths** have occurred among confirmed cases, **nearly half (46%)** of whom were **co-infected with HIV**.
- Disproportionate deaths in males and females** An equal number of deaths occurred among males and females (25 each). Among the male deaths, **72% (18/25)** were below 30 years of age, whereas among the female deaths, **68% (17/25)** were aged above 30 years.

### Weekly and Cumulative Confirmed Mpox Cases<sup>1</sup>

Year	Epi Week	Weekly Confirmed Cases	Cumulative Confirmed Cases	Percent Change (%)
2025	28	108	7530	1.46
2025	29	74	7604	0.98
2025	30	124	7728	1.63
2025	31	83	7811	1.07
2025	32	71	7882	0.91
2025	33	13	7895	0.16
2025	34	32	7927	0.41
2025	35	85	8012	1.07
2025	36	26	8038	0.32
2025	37	3	8041	0.04

<sup>1</sup> This is data for the last 10 Epiweeks. % Week-on-Week Change

## 2 Surveillance and Epidemiology


### 2.1 Summary Table of Incident Cases by Age and Sex

Cases by Age Group and Gender							
Distribution across demographics							
	Age Group	Total Cases	% of Cases <sup>1</sup>	Males	% of All Males <sup>2</sup>	Females	% of All Females <sup>2</sup>
	0-4	352	4.6%	202	4.6%	150	4.6%
	5-9	329	4.3%	185	4.2%	144	4.5%
	10-14	232	3.1%	125	2.9%	107	3.3%
	15-19	708	9.3%	335	7.7%	373	11.6%
	20-24	1,264	16.6%	561	12.8%	703	21.8%
	25-29	1,583	20.8%	867	19.9%	716	22.2%
	30-34	1,184	15.6%	722	16.5%	462	14.3%
	35-39	906	11.9%	612	14.0%	294	9.1%
	40-44	519	6.8%	364	8.3%	155	4.8%
	45-49	310	4.1%	230	5.3%	80	2.5%
	50+	206	2.7%	164	3.8%	42	1.3%
sum	—	7,593 <sup>3</sup>	100.0%	4,367	100.0%	3,226	100.0%
<sup>1</sup> Percentage of all cases within each age group							
<sup>2</sup> Percentage of all cases within each gender category							
<sup>3</sup> Total cases less due to some missing age information							


### 2.2 Summary Table of Incident Cases by the Top 10 Districts

Top 10 Districts Summary					
District	Total Cases	Total Deaths	Cases Last Epiweek	Deaths Last Epiweek	% Change in Cases <sup>1</sup>
Kampala District	2,721	16	0	0	–100.0%
Wakiso District	956	8	0	0	NA
Mbarara City	703	5	2	0	–33.3%
Masaka City	354	3	0	0	NA
Mukono District	290	2	0	0	–100.0%
Hoima City	218	2	2	0	–60.0%
Lyantonde District	143	0	0	0	NA
Nakasongola District	128	1	0	0	NA
Luwero District	112	0	0	0	NA
Mubende District	109	0	0	0	NA
<sup>1</sup> The percentage change in cases is calculated by comparing the number of cases in the most recent epidemiological week to the previous epidemiological week.					

### 2.3 Proportion of Affected Districts




## 2.4 Overall Distribution of Incident Cases by Time



## 2.5 Epicurves of the Top 10 Districts

### Weekly Epicurves of Top 10 Districts


*Note: Each panel uses a different y-axis scale to enhance readability*



## 2.6 Overall Distribution of Incident Cases by Time


## 2.7 Weekly Cumulative Cases

### Weekly Incident Mpox Cases & Cumulative (Confirmed Only)




## 2.8 Overall Distribution of Incident Cases by Age and Sex

Gender Distribution




Distribution of Age and Sex




## 2.9 Overall Distribution of Deaths by Place and Sex - Age

Mpox Deaths by District




Mpox Deaths by Age and Sex




## 2.10 Distribution of Deaths by HIV Status


Proportion of Mpox Deaths with HIV Comorbidity



Mpox Deaths by Age and Sex (HIV-related)



2.11 Overall Attack Rates by Age





2.12 Distribution of Incident Cases by Place(Absolute Numbers)



2.13 Attack Rates per 100K Population by Place



2.14 Attack Rates per 100K Population by Place and Top 10 Districts





# MPOX CASE DEFINITION

## COMMUNITY CASE DEFINITION

- An illness with signs of skin rash, fever, headache, muscle aches, back pain, and body weakness.

## SUSPECTED CASE

- Any person with skin rash on any part of the body not explained by any locally relevant common cause of papular or vesicular rash, mucosal lesions or swollen lymph nodes with acute onset of fever ( $>38.5^{\circ}\text{C}$ ), headache, myalgia (muscle pain/body ache), back pain, profound weakness, or fatigue.

## PROBABLE CASE

- A suspect case with history of contact to probable or confirmed Mpox case 21 days before the onset of signs or symptoms

## OR

- Has an epidemiological link to a probable or confirmed case of Mpox in the 21 days before symptom onset including multiple and/or casual sexual partners in the 21 days before symptom onset: has a positive test result for orthopoxviral infection (e.g., OPXV-specific PCR without MPXV-specific PCR or sequencing)



## CONFIRMED CASE

- A person with laboratory confirmed MPXV infection by detection of Unique sequences of Viral DNA by real-time polymerase chain reaction (RT-PCR) and/or sequence

## Contact Us

**Twitter:** @MinofHealthUG  
**Website:** <https://health.go.ug/>  
**Talk to Us:** 0800-100-066  
**Write to Us:** [helpdesk@health.go.ug](mailto:helpdesk@health.go.ug)

