GYTS Objectives

The Global Youth Tobacco Survey (GYTS), a component of the Global Tobacco Surveillance System (GTSS), is a global standard for systematically monitoring youth tobacco use (smoking and smokeless) and tracking key tobacco control indicators.

GYTS is a nationally representative school-based survey of students in grades associated with age 13 to 15 years and is designed to produce cross-sectional estimates for each country. GYTS uses a standard core questionnaire, sample design, and data collection protocol. It assists countries in fulfilling their obligations under the World Health Organization (WHO) Framework Convention on Tobacco Control (FCTC) to generate comparable data within and across countries. WHO has developed MPOWER, a package of selected demand reduction measures contained in the WHO FCTC:

mpower

Monitor tobacco use & prevention policies

Protect people from tobacco smoke

Offer help to quit tobacco use

Warn about the dangers of tobacco

Enforce bans on tobacco advertising, promotion, & sponsorship

Raise taxes on tobacco

GYTS Methodology

GYTS uses a global standardized methodology that includes a two-stage sample design with schools selected with a probability proportional to enrollment size. The classrooms within selected schools are chosen randomly and all students in selected classes are invited to participate in the survey. The survey uses a standard global core questionnaire with a set of optional questions that permits adaptation to meet the needs of the country on tobacco use and key tobacco control indicators. The questionnaire consists of the following topics: tobacco use (smoking and smokeless), cessation, secondhand smoke (SHS), pro- and anti-tobacco media and advertising, access and availability to obtain tobacco. The questionnaire is selfadministered; using scannable, paper-based bubble sheets, it is anonymous to ensure confidentiality.

In Mozambique, GYTS was conducted in 2013 by the Mozambique National Commission for UNESCO, under the coordination of the Ministry of Health. A total of 5,599 eligible students in grades 8-10 completed the survey, of which 3,062 were aged 13-15 years. The overall response rate of all students surveyed was 74.9%.





GYTS Highlights

TOBACCO USE

- 9.1% overall, 9.3% of boys, and 8.2% of girls currently used any tobacco products.
- 6.1% overall, 5.5% of boys, and 6.2% of girls currently smoked tobacco.
- 2.3% overall, 2.1% of boys, and 2.3% of girls currently smoked cigarettes.
- 4.3% overall, 5.0% of boys, and 3.3% of girls currently used smokeless tobacco.

SECONDHAND SMOKE

- 19.1% of students were exposed to tobacco smoke at home.
- 37.4% of students were exposed to tobacco smoke inside enclosed public places.

MEDIA

- Almost 6 in 10 students noticed anti-tobacco messages in the media.
- More than 3 in 10 students noticed tobacco advertisements or promotions when visiting points of sale.
- More than 1 in 10 students own something with a tobacco brand logo on it.

KNOWLEDGE & ATTITUDES

- 30.3% of students definitely thought other people's tobacco smoking is harmful to them.
- 70.8% of students favor banning smoking inside enclosed public places.






GYTS GLOBAL YOUTH TOBACCO SURVEY

FACT SHEET MOZAMBIQUE 2013

TOBACCO USE

SMOKED TOBACCO	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco smokers ¹	6.1	5.5	6.2
Current cigarette smokers ²	2.3	2.1	2.3
Frequent cigarette smokers ³	0.2	0.4	0.1
Current smokers of other tobacco ⁴	4.4	4.1	4.4
Ever tobacco smokers ⁵	15.4	16.0	14.3
Ever cigarette smokers ⁶	8.6	10.1	6.6
Ever smokers of other tobacco ⁷	9.1	8.2	9.6
SMOKELESS TOBACCO			
Current smokeless tobacco users ⁸	4.3	5.0	3.3
Ever smokeless tobacco users ⁹	8.8	9.4	7.9
TOBACCO USE (smoked and/or smokeless)			
Current tobacco users ¹⁰	9.1	9.3	8.2
Ever tobacco users ¹¹	20.0	20.6	18.6
SUSCEPTIBILITY			
Never to bacco users susceptible to to bacco use in the $\ensuremath{future^{12}}$	32.4	33.8	31.1
Never smokers who thought they might enjoy smoking a cigarette ¹³	6.6	6.9	6.3



CESSATION

	OVERALL (%)	BOYS (%)	GIRLS (%)
Current smokers who have ever received help/advice from a program or professional to stop smoking	21.7		

SECONDHAND SMOKE

	OVERALL (%)	BOYS (%)	GIRLS (%)
Exposure to tobacco smoke at home ⁺⁺	19.1	18.9	18.8
Exposure to tobacco smoke inside any enclosed public place ^{††}	37.4	35.3	39.3
Exposure to tobacco smoke at any outdoor public place ^{††}	38.7	38.0	39.6
Students who saw anyone smoking inside the school building or outside on school property [†]	21.4	22.3	20.5

MEDIA

TOBACCO ADVERTISING	OVERALL (%)	BOYS (%)	GIRLS (%)
	OVERALL (%)	B013 (%)	GIKLS (70)
Noticing tobacco advertisements or promotions at points of sale ¹⁴	34.7	31.1	37.9
Students who saw anyone using tobacco on television, videos, or movies ¹⁵	69.4	67.9	70.9
Students who were ever offered a free tobacco product from a tobacco company representative	7.1	7.7	6.5
Students who own something with a tobacco brand logo on it	12.4	13.1	11.6
ANTI-TOBACCO ADVERTISING			
Noticing anti-tobacco messages in the media [†]	58.3	57.4	59.2
Noticing anti-tobacco messages at sporting or community events ¹⁶	39.2	38.1	40.2
Current smokers who thought about quitting because of a warning label ¹⁷	41.8	38.4	46.9
Students who were taught in school about the dangers of tobacco use in the past 12 months	45.6	42.4	48.4

KNOWLEDGE & ATTITUDES

	OVERALL (%)	BOYS (%)	GIRLS (%)
Students who definitely thought it is difficult to quit once someone starts smoking tobacco	21.8	22.3	21.6
Students who thought smoking tobacco helps people feel more comfortable at celebrations, parties, and social gatherings	22.6	23.6	21.9
Students who definitely thought other people's tobacco smoking is harmful to them	30.3	29.6	31.0
Students who favor banning smoking inside enclosed public places	70.8	68.6	72.6
Students who favor banning smoking at outdoor public places	57.9	55.8	59.8

¹ Smoked tobacco anytime during the past 30 days. ² Smoked cigarettes anytime during the past 30 days. ³ Smoked cigarettes on 20 or more days of the past 30 days. ⁴ Smoked tobacco other than cigarettes anytime during the past 30 days. ⁵ Ever smoked any tobacco, even one or two puffs. ⁶ Ever smoked cigarettes, even one or two puffs. ⁷ Ever smoked tobacco other than cigarettes, even one or two puffs. ⁷ Ever smoked tobacco other than cigarettes, even one or two puffs. ⁷ Ever smoked tobacco other than cigarettes, even one or two puffs. ⁸ Used smokeless tobacco anytime during the past 30 days. ⁹ Ever used smokeless tobacco. ¹⁰ Smoked tobacco and/or used smokeless tobacco anytime during the past 30 days. ¹¹ Ever smoked tobacco and/or used smokeless tobacco. ¹² Susceptible to future tobacco use includes those who answered "Definitely yes", "Probably yes", or "Probably not" to using tobacco during the next 12 months. ¹³ Those who answered "Sefficiently yes", or "Probably not" to using tobacco during the next 12 months. ¹³ Those who answered "agree" or "Strongly agree" to the statement: "I think I might enjoy smoking a cigaretter. ¹⁴ Among those who visited a point of sale in the past 30 days. ¹⁵ Among those who matched television, videos, or movies in the past 30 days. ¹⁶ Among those who attended sporting or community events in the past 30 days. ¹⁷ Among those who noticed warning labels on cigarette packages in the past 30 days. ¹⁸ 2007 data is from Maputo City only. ¹ During the past 30 days. ¹⁰ During the past 7 days.

NOTE: Students refer to persons aged 13-15 years who are enrolled in school. Data have been weighted to be nationally representative of all students aged 13-15 years. Percentages reflect the prevalence of each indicator in each group, not the distribution across groups.

--Estimates based on unweighted cases less than 35 are suppressed.